C.O.W.S.® PROGRAM OVERVIEW

C.O.W.S.® Objectives: The C.O.W.S.® Program provides feedback to producers on cow comfort and management practices on their farm relative to regional benchmarks across the United States.

The ultimate goal is to engage the producer and management team to implement changes toward removing limitations on milk production and improving animal well-being.

The C.O.W.S.® Assessment Process

STEP 1: Herd Registration

Your herd is entered as a candidate herd by your NOVUS Sales Manager.

Herd qualifications:

- Requires animal restraint to attach loggers. Acceptable restraints include:
 - Parlors
- Tie-stalls
- Stanchions Headlocks
- Has a pen containing at least 40 lactating animals.
- Management team is eager to make changes toward improvement.

Complete herd information must be obtained:

- Consultant name and contact information.
- Farm name, contact person and physical address.
- Milking time for the assessment group.

STEP 2: Assessment Scheduling

Consultant and herd will be notified to schedule assessment.

STEP 3: On-Farm Assessment

Each assessment involves 2 visits to the dairy.

- Data loggers, which measure lying behavior, will be attached to 40 cows from the assessment group during milking (Figure 1).
- · Hock and knee health will be assessed at the same time.

FIGURE 1. Data Logger on a Cow's Leg



- Facility design (stall dimensions, bedding dry matter, etc.) and management (stocking density, bunk space, etc.) are evaluated before or after milking.
- · Management questions to be answered by the farm manager or herdsperson (30 - 45 minutes).

Visit 2:

- Cows will be gait scored as they exit the parlor.
- Logger removal will be done by the farm or nutritionist 4 to 5 days later and shipped back with a prepared, pre-paid shipping label.

STEP 4: Report Delivery

Report delivery will be scheduled with the producer and nutritionist when loggers are received and the report is ready. Turnaround time on the report varies between 1 to 3 weeks.

Once the results are finalized, they will be presented in a customized report. The management team will discuss opportunities for improvement and determine action plans with executable tasks.





C.O.W.S.® PROGRAM OVERVIEW



Research



- Support hoof health and structural integrity^{1,2}
- Improve reproductive performance^{1,2}
- Strengthen immune response^{2,3}

MHA MFP alimet.

- Maximize milk fat production by up to 0.3%^{4,5}
- Minimize the risk for diet-induced milk fat depression^{4,5}
- Increase rumen microbial protein production⁶

C.O.W.S.® Senior Commercial Manager Clem Nash, Ph.D.

P: 416.575.5882

E: Clemence.Nash@novusint.com

NORTHEAST, UPPER MIDWEST

Kelsey Hefter

C.O.W.S.® Technical Ruminant Specialist, NOVUS

P: 636.342.0989

E: Kelsey.Hefter@novusint.com

CANADA

Kaitlyn Dancy

Senior Sales Specialist, NOVUS

P: 587.436.7004

E: Kaitlyn.Dancy@novusint.com

References

- 1 Bach, A., A.M. Pinto and M. Blanch. 2015. Association between chelated trace mineral supplementation and milk yield, reproductive performance, and lameness in dairy cattle. Livestock Sci. 182: 69-75.
- 2 Zhao, X., Z.P. Li, J.H. Wang, X.M. Xing, Z.Y. Wang, L. Wang and Z.H. Wang. 2015. Effects of chelated Zn/Cu/Mn on redox status, immune responses and hoof health in lactating Holstein cows. J. Vet. Sci. 16(4): 439-446.
- 3 Nemec, L.M., J.D. Richards, C.A. Atwell, D.E. Diaz, G.I. Zanton, and T.F. Gressley, 2012. Immune responses in lactating Holstein cows supplemented with Cu, Mn, and Zn as sulfates or methionine hydroxy analogue chelates. J. DairySci. 95: 4568–4577.
- 4 Baldin, M., G. I. Zanton, and K. J. Harvatine. 2018. Effect of 2-hydroxy-4-(methylthio)butanoate (HMTBa) on risk of biohydrogenation-induced milk fat depression. J. Dairy. Sci. 101:376-385. https://doi.org/10.3168/jds.2017-13446
- 5 Baldin, M., H.A. Tucker, and K.J. Harvatine. 2019. Milk fat response and milk fat and urine marker prediction of biomarkers of microbial nitrogen flow during supplementation with 2-hydroxy-4-(methylthio)butanoate (HMTBa). J. Dairy Sci. 102:6157-6166. HYPERLINK "https://linkinghub.elsevier.com/retrieve/pii/S0022030219303935"DOI: 10.3168/jds.2018-15031
- 6 Pitta, D.W., N. Indugu, B. Vecchiarelli, M. Hennessy, M. Baldin, and K.J. Harvatine. 2020. Effect of 2-hydroxy-4-(methylthio) butanoate (HMTBa) supplementation on rumen bacterial populations in dairy cows when exposed to diets with risk for milk fat depression. J. Dairy Sci. 103:2718-2730. HYPERLINK "https://www.journalofdairyscience.org/article/S0022-0302[19]31110-5/fulltext"DOI: 10.3168/jds.2019-17389

Novus International, Inc. | 17988 Edison Ave. | Chesterfield, MO 63005 USA | 1-800-568-0088 | novusint.com

Products not available in all countries.

NOTICE: While the information contained herein ("Information") is presented in good faith and believed to be correct as of the date hereof, Novus International, Inc., does not guarantee satisfactory results from reliance upon such information, disclaims all liability for any loss or damage arising out of any use of this information or the products to which said information refers and MAKES NO REPRESENTATIONS OR WARRANTIES, EITHER EXPRESS OR IMPLIED, OF MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE OR OF ANY OTHER NATURE WITH RESPECT TO THE INFORMATION OR PRODUCTS, except as set forth in NOVUS's standard conditions of sale. Nothing contained herein is to be construed as a recommendation to use any product or process in conflict with any patent, and Novus International, Inc. makes no representation or warranty, express or implied, that the use thereof will not infringe any patent.

